



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBF180Mu01 100µg

Recombinant 17-Beta-Hydroxysteroid Dehydrogenase Type 10 (HSD17b10)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ala4~Ala202

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: O08756

Host: *E. coli*

Subcellular Location: Mitochondrion.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.


Predicted isoelectric point: 8.9 Predicted Molecular Mass: 24.1kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

**AVRSVKG LVAVVTGGAS GPWLATAKRL VGQGATAVLL DVPDSEGESQ AKKLGESCIF
APANVTSEKE IQAALTLAKE KFGRIDVAVN CAGIAVAIKT YHQKKNKIHT LEDFQRVINV
NLIGTFNVIR LVAGEMGQNE PDQGGQRGVI INTASVAAFE GQVGQAAYSA SKGGIDGMTL
PIARDLPTG IRVVTIAPGL FA**