TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB421Hu01 100µg Recombinant Nucleoporin 210kDa (NUP210) Organism Species: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Leu1288~Val1449 (Accession # Q8TEM1), with two kDa 70 N-terminal Tags, His-tag and T7-tag. 44 Host: E. coli 33 Subcellular Location: Nucleusnuclear pore 26 complex. Nucleus membrane; Single-pass type I 22 membrane protein. Endoplasmic reticulum membrane. 18 **Purity: >95%** Endotoxin Level: <1.0EU per 1µg 14 (determined by the LAL method). 10 Formulation: Supplied as lyophilized form in PBS, pH7.4, 15% SDS-PAGE containing 5% trehalose, 0.01% sarcosyl. Predicted isoelectric point: 6.5 Predicted Molecular Mass: 21.3kDa Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-LLN PEIEAEQILM SPNSYIKLQT NRDGAASLSY RVLDGPEKVP VVHVDEKGFL ASGSMIGTST IEVIAQEPFG ANQTIIVAVK VSPVSYLRVS MSPVLHTQNK EALVAVPLGM TVTFTVHFHD NSGDVFHAHS SVLNFATNRD DFVQIGKGPT NNTCVVRTV