TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBF173Hu01 100ug

Recombinant 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3)

Organism Species: Homo sapiens (Human)

Instruction manual

kDa 70

14

10

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Met1~Arg310

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: P37058

Host: E. coli

Subcellular Location: Endoplasmic reticulum membrane.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL

method).

Formulation: Supplied as lyophilized form in 20mM Tris, 15% SDS-PAGE

150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 8.9

Predicted Molecular Mass: 38.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

MGDVLEQFFI LTGLLVCLAC LAKCVRFSRC VLLNYWKVLP KSFLRSMGQW AVITGAGDGI GKAYSFELAK RGLNVVLISR TLEKLEAIAT EIERTTGRSV KIIQADFTKD DIYEHIKEKL AGLEIGILVN NVGMLPNLLP SHFLNAPDEI QSLIHCNITS VVKMTQLILK HMESRQKGLI LNISSGIALF PWPLYSMYSA SKAFVCAFSK ALQEEYKAKE VIIQVLTPYA VSTAMTKYLN TNVITKTADE FVKESLNYVT IGGETCGCLA HEILAGFLSL IPAWAFYSGA FQRLLLTHYV AYLKLNTKVR