

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA635Hu01 100µg **Recombinant Complement Factor H (CFH)** Organism Species: Homo sapiens (Human) Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

#### [PROPERTIES]

Residues: Ser860~Arg1231 Tags: N-terminal His-Tag Accession: P08603 Host: E. coli Subcellular Location: Secreted. **Purity: >95%** Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). 15% SDS-PAGE Formulation: Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. Predicted isoelectric point: 7.1 Predicted Molecular Mass: 43.2kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

# [USAGE]

Reconstitute in ddH<sub>2</sub>O.



### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

# [<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

S IPLCVEKIPC SQPPQIEHGT INSSRSSQES YAHGTKLSYT CEGGFRISEE NETTCYMGKW SSPPQCEGLP CKSPPEISHG VVAHMSDSYQ YGEEVTYKCF EGFGIDGPAI AKCLGEKWSH PPSCIKTDCL SLPSFENAIP MGEKKDVYKA GEQVTYTCAT YYKMDGASNV TCINSRWTGR PTCRDTSCVNPPTVQNAYIVSRQMSKYPSGERVRYQCRSPYEMFGDEEVM CLNGNWTEPP QCKDSTGKCG PPPPIDNGDI TSFPLSVYAP ASSVEYQCQNLYQLEGNKRI TCRNGQWSEP PKCLHPCVIS REIMENYNIA LRWTAKQKLY SRTGESVEFV CKRGYRLSSR SHTLRTTCWD GKLEYPTCAK R

## [REFERENCES]

- 1. Ripoche J., et al. (1988) Biochem. J. 249:593-602.
- 2. Kristensen T., et al. (1986) J. Immunol. 136:3407-3411.
- 3. Estaller C., et al. (1991) J. Immunol. 146:3190-3196.
- 4. Fenaille F., et al. (2007) Glycobiology 17:932-944.