TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA798Ga02 100µg

Recombinant Hypoxia Inducible Factor 1 Alpha (HIF1a)

Organism Species: Chicken (Gallus)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Asp238~Lys389 (Accession # Q9YIB9), with

N-terminal His-Tag.

Host: E. coli

Subcellular Location: Cytoplasm. Nucleus.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg

(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris,

500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 9.3


Predicted Molecular Mass: 18.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in ddH₂O.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGS-DSK TFLSRHSLDM KFSYCDERIT ELMGYEPEEL LGRSIYEYYH ALDSDHLTKT HHDMFTKGQV TTGQYRMLAK QGGYVWVETQ ATVIYNTKNS QPQCIVCVNY VLSGIVQKDL IFSLGQTECM LKPVESPEMK MTKIFSKDDW DDTNSLFEK

[REFERENCES]

- 1. Takahashi T. (1998) Submitted to the EMBL/GenBank/DDBJ databases.
- 2. Ratcliffe PJ. (2003) Blood Purif. 20 (5): 445-50.
- 3. van Uden P., et al. (2008) Biochem J. 412 (3): 477-484.
- 4. Cho, S; et al. (2001) FEBS Lett. (Netherlands) 498 (1): 62-6.