



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA413Mu01 10 $\mu$ g

Recombinant Mucin 1 (MUC1)

Organism Species: *Mus musculus* (Mouse)

*Instruction  
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul,  
2013)

## [ PROPERTIES ]

Residues: Ser410~Ala525 (Accession # Q02496),  
with N-terminal His-Tag.


Host: *E. coli*

Subcellular Location: Apical cell membrane; Single-pass  
type I membrane protein. Cell membrane. Cytoplasm.  
Nucleus.

Purity: >95%

Endotoxin Level: <1.0EU per 1 $\mu$ g (determined by the  
LAL method).

Formulation: Supplied as lyophilized form in 20mL  
Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM  
DTT, 0.01% sarcosyl, 5% trehalose, and preservative.



The possible reasons that the actual band size differs from the predicted are as follows:

Predicted isoelectric point:

6.0 Predicted Molecular Mass:

14.6kDa



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Accurate Molecular Mass: 10kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

## [ USAGE ]

Reconstitute in sterile PBS, pH7.2-pH7.4.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCES ]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHSGSEF-S PQLSVGVSFF FLSFYIQNHP FNSSLED PSS NYYQELKRNI



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

SGLFLQIFNG DFLGISSIKF RSGSVVVEST VVFREGTFSA SDVKSQLIQH KKEADDYNLT  
ISEVKVNEMQ FPPSA

## [ REFERENCES ]

1. Spicer A.P., *et al.* (1991) J. Biol. Chem. 266:15099-15109.
2. Vos H.L., *et al.* (1991) Biochem. Biophys. Res. Commun. 181:121-130.
3. McAuley J.L., *et al.* (2007) J. Clin. Invest. 117:2313-2324.
4. VanLith M.L., *et al.* (2002) Int. Immunol. 14:873-882.