



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

**YBC903Ra01 10 $\mu$ g**

**Recombinant Deiodinase, Iodothyronine, Type II (DIO2)**

**Organism Species: Rattus norvegicus (Rat)**

***Instruction manual***

**FOR IN VITRO USE AND RESEARCH USE ONLY**

**NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES**

10th Edition (Revised in Jan, 2014)

## **[ PROPERTIES ]**

**Residues: Ser30~Phe259**

**Tags: Two N-terminal Tags, His-tag and MBP-tag**

**Accession: P70551**

**Host: *E. coli***

**Subcellular Location: Membrane; Single-pass membrane protein.**

**Purity: >95%**

**Endotoxin Level: <1.0EU per 1 $\mu$ g (determined by the LAL method).**

**Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative.**

**Predicted isoelectric point: 5.8**

**Predicted Molecular Mass: 72.4kDa**

**Accurate Molecular Mass: 60kDa as determined by SDS-PAGE reducing conditions.**


**Applications: SDS-PAGE; WB; ELISA; IP.**

**(May be suitable for use in other assays to be determined by the end user.)**

The possible reasons that the actual band size differs from the predicted are as follows:

### **Note:**

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.





## **[ USAGE ]**

Reconstitute in sterile PBS, pH7.2-pH7.4.

## **[ STORAGE AND STABILITY ]**

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCES ]**

The sequence of the target protein is listed below.

S VILLKHVALL LSRSKSTRGE WRRMLTSEGL RCVWNSFLLD AYKQVKLGED APNSSVVHVS  
NPEAGNNCAS EKTADGAECH LLDFAAERP LVVNFGSATU PPFTRQLPAF RQLVEEFSSV  
ADFLLVYIDE AHPDGDWAVP GDSSMSFEVK KHRNQEDRCA AAHQLLERFS LPPQCQVVAD  
RMDNNANVAY GVAFERVCIV QRRKIAYLGG KGPFSYNLQE VRSWLEKNF

## **[ REFERENCES ]**

1. Croteau W., *et al.* (1996) J. Clin. Invest. 98:405-417.
2. Gondou A., *et al.* (1998) Thyroid 8:615-622.
3. Murakami M., *et al.* (1997) Neurosci. Lett. 227:65-67.
4. Bailey M.J., *et al.* (2009) J. Biol. Chem. 284:7606-7622.