TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB386Mu01 100µg

Recombinant B-Lymphocyte Activation Antigen B7-1 (LAB7-1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

70 🐷

[PROPERTIES]

Residues: Asp39~Asn246

Tags: Two N-terminal Tags, His-tag and GST-tag

Accession: Q00609

Host: E. coli

Subcellular Location: Membrane; Single-pass

type I membrane protein.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the

LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 6.3

Predicted Molecular Mass: 53.7kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



Web:www.ybio.net Email:shybio@126.com

Reconstitute in sterile PBS, pH7.2-pH7.4.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

DE QLS K S V K D K V LL P C RY N S P H ED E S E D R I Y W QK H D K V V L S V IA G K LK V W P E YKNRTLYDNT TYSLIILGLV LSDRGTYSCV VQKKERGTYE VKHLALVKLS IKADFSTPNI TESGNPSADT KRITCFASGG FPKPRFSWLE NGRELPGINT TISQDPESELYTISSQLDFN TTRNHTIKCL IKYGDAHVSE DFTWEKPPED PPDSKN