



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA149Hu01 10 μ g
Collagen Type IV Alpha 1 (COL4a1)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: His1444~Thr1669 (Accession # P02462), with two N-terminal Tags, His-tag and S-tag.

Host: *E. coli*

Subcellular Location: Secreted, extracellular space, extracellular matrix, basement membrane.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 6.7 Predicted


Molecular Mass: 30.7kDa Applications:

SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8^oC for one month.

Aliquot and store at -80^oC for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37^oC for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and S-tag, its sequence is listed below.

MHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF-
HGFLVTR HSQTIDDPQC PSGTKILYHG YSLLYVQGNE RAHGQDLGTA GSCLRKFSTM
PFLFCNINNV CNFASRNDYS YWLSTPEPMP MSMAPITGEN IRPFISRCV CEAPAMVMAV
HSQTIQIPPC PSGWSSLWIG YSFVMHTSAG AEGSGQALAS PGSCLEEFRS APFIECHGRG
TCNYANAYS FWLATIERSE MFKKPTPSTL KAGELRTHVS RCQVCMRRT

[REFERENCES]

1. Soininen R., *et al.* (1988) FEBS Lett 225 (1-2): 188–94.
2. Plaisier E., *et al.* (2007) N. Engl. J. Med. 357 (26): 2687–2695.
3. Tryggvason K., *et al.* (1990) Ann. N. Y. Acad. Sci. 580: 97–111.
4. Hinek A. (1995) Cell Adhes. Commun. 2 (3): 185–93.
5. Stander M., *et al.* (1999) Cell Tissue Res. 296 (2): 221–7.