



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB972Mu01

50µg

Recombinant Pigment Epithelium Derived Factor (PEDF)

Organism Species: Mus musculus (Mouse)

***Instruction
manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul,
2013)

[PROPERTIES]

Residues: Ser19~Leu389 (Accession # P97298), with two N-terminal Tags, His-tag and S-tag.

Host: *E. coli*

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative.

Predicted isoelectric point: 6.2

Predicted Molecular Mass: 47.0kDa

Accurate Molecular Mass: 55kDa as determined by SDS-PAGE reducing


The possible reasons that the actual band size differs from the predicted are as follows:
conditions.

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.

5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and S-tag, its sequence is listed below.

MHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF- SQ
NVPSSSEGSP VPDSTGEPVE EEDPFFKVPV NKLAAAVSNF GYDLYRLRSS ASPTGNVLLS
PLSVATALSA LSLGAHRTE SVIHRALYYD LITNPDIHST YKELLASVTA PEKNLKSASR
IVFERKLRVK SSFVAPLEKS YGTRPRILTG NPRVDLQEIN NWWQAQMKGK IARSTREMP
ALSILLGLVA YFKGQWVTKF DSRKTTLQDF HLEDRTVRVPMMSDPKAIL RYGLDSDLNC
KIAQLPLTGS MSIIFLPLT VTQNLTMIEE SLTSEFIHDI DRELKTIQAV LTVPKLKLSF
EGELTKSLQD MKLQSLFESP DFSKITGKPV KLTQVEHRAA FEWNEEGAGS SPSPGLQPVR
LTFPLDYHL

[REFERENCES]

1. Shirozu M., *et al.* (1996) Genomics 37:273-280.
2. Kozaki K., *et al.* (1998) J. Biol. Chem. 273:15125-15130.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

3. Singh V.K., *et al.* (1998) Mol. Vis. 4:7-7.

4. Abramson L.P., *et al.* (2003) J. Pediatr. Surg. 38:336-342.