



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

**YBB690Hu01 100μg**

**Recombinant Permeability Glycoprotein (Pgp)**

**Organism Species: Homo sapiens (Human)**

*Instruction  
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May,  
2016)

## [ PROPERTIES ]

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Phe394~Val672

**Tags:** Two N-terminal Tags, His-tag and

GST-tag **Tissue Specificity:** Liver, Kidney,

Brain.

**Subcellular Location:** Cell membrane; Multi-pass membrane protein.

**Purity:** >98%

**Traits:** Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

**Original Concentration:** 200ug/mL

**Applications:** SDS-PAGE; WB; ELISA; IP; CoIP; ReporterAssays;

Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

**Predicted isoelectric point:**

**7.0 Predicted Molecular Mass:**

57.9kDa

**Accurate Molecular Mass:** 61kDa as determined by SDS-PAGE reducing conditions.

**Phenomenon explanation:**

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

## [ USAGE ]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCE ]



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

FRNVHFS  
YPSRKEVKIL KGLNLKVQSG QTVALVGNSG CGKSTTVQLM QRLYDPTEGM  
VSVDGQDIRT INVRFLREII GVVSQEPVLF ATTIAENIRY GRENTMDEI  
EKAVKEANAY DFIMKLPHKF DTLVGERGAQ LSGGQKQRIA IARALVRNPK  
ILLLDEATSA LDTSEAVVQ VALDKARKGR TTIVIAHRLS TVRNADVIAG  
FDDGVIVEKG NHDELMKEKG IYFKLVTMQT AGNEVELENA ADESKSEIDA  
LEMSSNDSRS SLIRKRSTRR SV



TEL:4006-871-

227 Web:www.ybio.net Email:sh

ybio@126.com

## [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE