TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBC178Hu01 100µg

Recombinant Neurocan (NCAN)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

> 10th Edition (Revised in Jan, 2014)

## [ PROPERTIES ]

Residues: His1008~Lys1273

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: 014594

Host: E. coli

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level:  $\langle 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per } 1 \mu \text{ g} \text{ (determined by } 1.0EU \text{ per }$ 

the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 5.6


The possible reasons that the actual band size differs from the predicted are as follows:

Predicted Molecular Mass: 33.6kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing

conditions.

Applications: SDS-PAGE; WB; ELISA; IP.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

(May be suitable for use in other assays to be determined by the end user.)

#### Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

### [ USAGE ]

Reconstitute in sterile PBS, pH7. 2-pH7. 4.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

# [ SEQUENCES ]

The sequence of the target protein is listed below.

HSD PCENNPCLHG GTCNANGTMY GCSCDQGFAG ENCEIDIDDC LCSPCENGGT

C I D E V N G F V C LC L P S Y G G S F C E K D T E G C D R G W H K F Q G H C Y RY FA H R R AW Е

DAEKDCRRRS GHLTSVHSPE EHSFINSFGH ENTWIGLNDR IVERDFQWTD NTGLQFENWR



ENQPDNFFAGGEDCVVMVAHESGRWNDVPCNYNLPYVCKKGTVLCGPP PA

VENASLIGAR KAKYNVHATV RYQCNEGFAQ HHVATIRCRS NGKWDRPQIV CTK