

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA143Si01 20µg

Recombinant Vascular Endothelial Growth Factor A (VEGFA)

Organism Species: Rhesus monkey (Simian)

Instruction manual

kDa 70

44 33

26

22 18

14

10

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Pro28 Arg147 (Accession #

NP 001265339.1), with N-terminal His-Tag.

Host: E. coli

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20ml 15% SDS-PAGE

Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM

DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 6.3

Predicted Molecular Mass: 15.5kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



Reconstitute in sterile PBS, pH7.2-pH7.4

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF-PMA EGGGQNHHEV VKFMDVYQRS YCHPIETLVD IFQEYPDEIE YIFKPSCVPL MRCGGCCNDE GLECVPTEES NITMQIMRIK PHQGQHIGEM SFLQHNKCEC RPKKDRARQE KCDKPRR

[REFERENCES]

- 1. Aschenbach L. C., et al. (2012) Contraception 0:0-0.
- 2. Mackenzie., et al. (2012) Development (n. d.): 1371-380.
- 3. Creuzet., et al. (2002) Development (n.d.): 4301-313.
- 4. Wu LW., et al. (2000) J. Biol. Chem. 275 (9): 6059 6062.