



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA653Hu01 100 μ g
Recombinant Bone Morphogenetic Protein 1 (BMP1)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Glu610~Ser843 (Accession # P13497), with two N-terminal Tags, His-tag and GST-tag.

Host: *E. coli*

Subcellular Location: Golgi apparatus, trans-Golgi network. Secreted, extracellular space, extracellular matrix.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g (determined by the LAL method).


Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative.

Predicted isoelectric point: 6.1

Predicted Molecular Mass: 58.5kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



[USAGE]



TEL:4006-871-

227 Web:www.ybio.net Email:shybio@126.com

Reconstitute in sterile PBS, pH7.2-pH7.4.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and GST-tag, its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID
GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV
DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK
KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPKSD GSTSGSHHH HHHSAGLVPR
GSTAIGMKET AAAKFERQHM DSPDLGTLEV LFQGPLGSEF-E YPPNKNCIWQ LVAPTQYRIS
LQFDFFETEG NDVCKYDFVE VRSGLTADSK LHGKFCGSEK PEVITSQYNN MRVEFKSDNT
VSKKGFKAHFFSDKDEC SKDN GG C Q Q DCVNTFGSYECQC RSGF VLHD NKH
DCKEAGCDHK VTSTSGTITS PNWPDKYPSK KECTWAISS PGHRVKLTFM EMDIESQPEC
AYDHLEVFDG RDAKAPVLGR FCGSKKPEPV LATGSRMFLR FYS

[REFERENCES]

1. Li S.W., et al. (1996) Proc. Natl. Acad. Sci. U.S.A. 93:5127-5130.
2. Takahara K., et al. (1994) J. Biol. Chem. 269:32572-32578.
3. Garrigue-Antar L., et al. (2001) J. Biol. Chem. 276:26237-26242.
4. Mac Sweeney A., et al. (2008) J. Mol. Biol. 384:228-239.