



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA126Ra01 100μg

Recombinant TEK Tyrosine Kinase, Endothelial (Tie2)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Cys44~Glu341

Tags: Two N-terminal Tags, His-tag and GST-tag

Accession: D3ZCD0

Host: *E. coli*

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 8.0


Predicted Molecular Mass: 63.6kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

**CIASGWH PHEPITIGRD FEALMNQHQD PLEVTQDVTR EWAKKVVWKR EKASKINGAY
FCEGRVRGQA IRIRTMKMRQ QASFLPATLT MTVDRGDNVN ISFKKVLKE EDAVIYKNGS
FIHSVPRHEV PDILEVHLPH AQPQDAGVYS ARYIGGNLFT SAFTRLIVRR CEAQKWGPDC
NRPCTTCKNNGVCHEDTGECICPPGFMGRTCEKACEPHTFGRTCKERC SG
SEGCKSYVFCLPDPYGCSCATGWRGLQCNEACPYGHYGPDC KLRCHCTNE
EMCDRFQGCL CSQGWQGLQC E**