

YBJ094Hu01 10µg

Recombinant Fructosamine-3-Kinase (FN3K)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Met1~Lys309 Tags: N-terminal His-Tag

Purity: >95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Reporter Assays; Purification;

Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.1

Predicted Molecular Mass: 38.9kDa

Accurate Molecular Mass: 40kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8_oC for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MEOLLRAELR TATLRAFGGP GAGCISEGRA YDTDAGPVFV KVNRRTOARO MFEGEVASLE ALRSTGLVRV PRPMKVIDLP GGGAAFVMEH LKMKSLSSQA SKLGEOMADL HLYNOKLREK LKEEENTVGR RGEGAEPQYV DKFGFHTVTC CGFIPOVNEW ODDWPTFFAR HRLOAOLDLI EKDYADREAR ELWSRLOVKI PDLFCGLEIV PALLHGDLWS GNVAEDDVGP IIYDPASFYG HSEFELAIAL MFGGFPRSFF TAYHRKIPKA PGFDORLLLY OLFNYLNHWN HFGREYRSPS LGTMRRLLK

[IDENTIFICATION]




Figure 1. SDS-PAGE



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com