

YBB843Hu01 100ug Recombinant Signaling Lymphocytic

Activation Molecule Family, Member 2 (SLAMF2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[<u>PROPERTIES</u>]


Residues: Gln27^{Thr243} Tags: Two N-terminal Tags, His-tag and T7-tag Accession: P09326 Host: *E. coli* Subcellular Location: Cell membrane. Lipid-anchor GPI-anchor . Purity: >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mN [Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mN [DTT, 0.01% sarcosyl, 5% trehalose, and preservative. Predicted isoelectric point: 8.7

_

Predicted Molecular Mass:

28.6kDa





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that incubate the protein at 37° C for 48h, and no obvious degradation and is. precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below. QGHL VHMTVVSGSN VTLNISESLP ENYKQLTWFY TFDQKIVEWD SRKSKYFESK FKGRVRLDPQ SGALYISKVQ KEDNSTYIMR VLKKTGNEQE WKIKLQVLDP VPKPVIKIEK IEDMDDNCYL KLSCVIPGES VNYTWYGDKR PFPKELQNSV LETTLMPHNY SRCYTCQVSN SVSSKNGTVC LSPPCTLARS FGVEWIASWL VVTVPTILGL LLT