



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB796Ra01 100μg

Recombinant Signal Transducer And Activator Of Transcription 2 (STAT2)

Organism Species: Rattus norvegicus (Rat)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Arg587~Ile838 (Accession # Q5XI26),
with two N-terminal Tags, His-tag and T7-tag.

Host: *E. coli*

Subcellular Location: Cytoplasm. Nucleus.

Purity: >95%

Endotoxin Level: <1.0EU per 1μg
(determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS,
pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 4.9


Predicted Molecular Mass: 32.7kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGS S HH H HH H SSG LVP R G SH MA SMT G GQ Q M GR G S EF- RR LL KK MVS G TFL L

RFSETSEGGI TCSWVEHQDD DKVLIYSVQP YTKEVLQSLP LTEIIRHYQV LAEENAPENP

LRFLYPRVPR DEAFGCYYQE KVNLEERRKY LKHRLIVISN RQADELQQPL ELKQDPESLE

LDAELMSVME LARDLELQSM LQVGLDLGVE PKPDPTLSPA PQILLEPDPA RALNQLLPEP

DLPQDLQQLN TGEMEIFRNN INIDDVMPYG DPVLAGQSII EEAYRSCPSH FDVDGPLI