

YBA728Hu01 5µg

Recombinant Prostaglandin Endoperoxide Synthase 1 (PTGS1) Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul. 2013)

[PROPERTIES]

Residues: Gly367~Leu599 (Accession # P23219), with

two N-terminal Tags, His-tag and T7-tag.

Host: E. coli

Subcellular Location: Microsome membrane;

Peripheral membrane protein. Endoplasmic

reticulum membrane.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the

LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM

DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 6.1

Predicted Molecular Mass: 30.4kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



227 Web:www.ybio.net Email:shybio@

126. com

Reconstitute in sterile PBS, pH7.2-pH7.4.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGSSHH HHHH SSGLVPRG SH MASMTG GQQ M GR GSEF- GVQF QYRNRIAMEF NHLYHWHPLM PDSFKVGSQE YSYEQFLFNT SMLVDYGVEA LVDAFSRQIA GRIGGGRNMD HHILHVAVDV IRESREMRLQ PFNEYRKRFG MKPYTSFQEL VGEKEMAAEL EELYGDIDAL EFYPGLLLEK CHPNSIFGES MIEIGAPFSL KGLLGNPICS PEYWKPSTFG GEVGFNIVKT ATLKKLVCLN TKTCPYVSFR VPDASQDDGP AVERPSTEL

[REFERENCES]

- 1. Yokoyama C., Tanabe T. (1989) Biochem. Biophys. Res. Commun. 165:888-894.
- 2. Funk C.D., et al. (1991) FASEB J. 5:2304-2312.
- 3. Takahashi Y., et al. (1992) Biochem. Biophys. Res. Commun. 182:433-438.
- 4. Diaz A., et al. (1992) J. Biol. Chem. 267:10816-10822.