



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA478Hu01 50μg

Recombinant Troponin I Type 3, Cardiac (TNNI3)

Organism Species: Homo sapiens (Human)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul,
2013)

[PROPERTIES]

Residues: Met1~Gly203 (Accession # P19429), with
N-terminal His-Tag.


Host: *E. coli*

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by the
LAL method).

Formulation: Supplied as lyophilized form in PBS,
pH7.4, containing 1mM DTT, 5% trehalose, 0.05%
sarcosyl and preservative.

Predicted isoelectric point: 9.7 Predicted



The possible reasons that the actual band size differs from the predicted are as follows:

Molecular Mass: 24.6kDa

Accurate Molecular Mass: 28kDa as determined by SDS-PAGE reducing
conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end
user.)



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHSGSEF- MADGSSDAAR EPRPAPAPIR RRSSNYRAYA TEPHAKKSK
ISASRKLQLK TLLQIAKQE LERAEERRG EKGRALSTRC QPLELAGLGF AELQDLQRQL
HARVDKVDEE RYDIEAKVTK NITEIADLTQ KIFDLRGKFK RPTLRRVRIS ADAMMQALLG
ARAKESLDLR AHLKQVKKED TEKENREVG DWRKNIDALSG MEG



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[REFERENCES]

1. Vallins W.J., *et al.* (1990) FEBS Lett. 270:57-61.
2. Armour K.L., *et al.* (1993) Gene 131:287-292.
3. Bhavsar P.K., *et al.* (1996) Genomics 35:11-23.
4. Mittmann K., *et al.* (1990) FEBS Lett. 273:41-45.