

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBD251Mu01 50µg Recombinant Ferritin, Mitochondrial (FTMT) Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PROPERTIES ]

Residues: Ser54~His229 (Accession # Q9D5H4), with

N-terminal His-Tag.

Host: E. coli

Subcellular Location: Mitochondrion.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris

150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01%

sarcosyl, 5% trehalose, and preservative.


Predicted isoelectric point: 6.1

Predicted Molecular Mass: 21.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

# [ USAGE ]





Reconstitute in ddH2O.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

### [ SEQUENCES ]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF- SSQDSTR PSRVRQNFHP DSEAAINRQI NLELYASYVY LSMAYYFSRD DVALYNFSKY FLRQSLEERE HAEKLMKLQN QRGGRICLQD IKKPDKDDWE CGLRAMECAL LLEKNVNQSL LDLHTLASEK GDPHLCDFLE THYLHEQVKS IKELGDHVHN LVTMGAPAAG LAEYLFDKH

#### [ REFERENCES ]

- 1. Nie G., et al. (2005) Blood 105:2161-2167.
- 2. Wilkinson J., et al. (2006) Biochem. J. 395:501-507.
- 3. Nie G., et al. (2006) Blood 108:2428-2434.
- 4. Iwasaki K., et al. (2007) J. Biol. Chem. 282:22335-22343.