



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

**YBE787Mu01 500μg**  
**Recombinant Annexin A1 (ANXA1)**  
**Organism Species: Mus musculus (Mouse)**  
***Instruction manual***

**FOR IN VITRO USE AND RESEARCH USE ONLY**  
**NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES**

9th Edition (Revised in Jul, 2013)

**[ PROPERTIES ]**

**Residues:** Glu6~Asn346 (Accession # P10107), with two N-terminal Tags, His-tag and S-tag.

**Host:** *E. coli*

**Subcellular Location:** Nucleus. Cytoplasm. Cell projection, cilium. Basolateral cell membrane.

**Purity:** >95%

**Endotoxin Level:** <1.0EU per 1μg (determined by the LAL method).

**Formulation:** Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 0.01% sarcosyl 5% trehalose, and preservative.

**Predicted isoelectric point:** 6.2

**Predicted Molecular Mass:** 43.9kDa

**Accurate Molecular Mass:** 40kDa as determined by SDS-PAGE reducing conditions.


**Applications:** SDS-PAGE; WB; ELISA; IP.

**(May be suitable for use in other assays to be determined by the end user.)**

The possible reasons that the actual band size differs from the predicted are as follows:

**Note:**

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.





## **[ USAGE ]**

Reconstitute in sterile PBS, pH7.2-pH7.4.

## **[ STORAGE AND STABILITY ]**

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCES ]**

The target protein is fused with two N-terminal Tags, His-tag and S-tag, its sequence is listed below.

MHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS- EFLKQ  
ARFLENQE QE YVQAVKSYKG GPGSAVSPYP SFNVSSDVAA LHKAIMVKGV DEATIIDILT  
KRTNAQRQQI KAAYLQENGK PLDEVLRKAL TGHLEEVVLA MLKTPAQFDA DELRGAMKGL  
GTDEDTLIEI LTTRSNEQIR EINRVYREEL KRDLAKDITS DTSGDFRKAL LALAKGDRCQ  
DLSVNQDLAD TDARALYEAG ERRKGTDVNV FTTILTSRSF PHLRRVFQNY GKYSQHDMNK  
ALDLELKGDI EKCLTTIVKC ATSTPAFFAE KLYEAMKGAG TRHKALIRIM VSRSEIDMNE  
IKVFYQKKYG ISLCQAILDE TKG DY EKILV ALCGGN

## **[ REFERENCES ]**

1. Philipps C., *et al.* (1989) Biochem. Biophys. Res. Commun. 159:155-162.
2. Lennon N.J., *et al.* (2003) J. Biol. Chem. 278:50466-50473.
3. Muimo R., *et al.* (2000) J. Biol. Chem. 275:36632-36636.
4. Hannon R., *et al.* (2003) FASEB J. 17:253-255.