



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA154Hu01 100μg

Recombinant Carbohydrate Antigen 125 (CA125)

Organism Species: Homo sapiens (Human)

*Instruction  
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan,  
2014)

## [ PROPERTIES ]

Residues: Arg21622~Ser21762

Tags: Two N-terminal Tags, His-tag and S-tag

Accession: Q8WXI7

Host: *E. coli*


Subcellular Location: Cell membrane; Single-pass type I  
membrane protein. Secreted, extracellular Space.

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by the  
LAL method).

Formulation: Supplied as lyophilized form in 20ml

Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,  
1mM DTT, 0.01% sarcosyl, 5% trehalose, and



The possible reasons that the actual band size differs from the predicted are as follows:  
preservative.

Predicted isoelectric point:

6.2 Predicted Molecular Mass:

21.2kDa



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Accurate Molecular Mass: 26kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

**Note:**

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[ USAGE ]

Reconstitute in ddH<sub>2</sub>O.

[ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[ SEQUENCES ]

The sequence of the target protein is listed below.

RVIALRSVK NGAETRVDLL CTYLQPLSGP GLPIKQVFHE LSQQTGTR LGPYSLDKDS



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

LYLNGYNEPG LDEPPTPKP ATTFLPPLSE ATTAMGYHLK TLTLNFTISN LQYSPDMGKG

SATFNSTEGV LQHLLRPLFQ KS

## [ REFERENCES ]

1. O'Brien T.J., *et al.* (2002) Tumor Biol. 23:154-169.
2. O'Brien T.J., *et al.* (2001) Tumor Biol. 22:348-366.
3. Yin B.W.T., Lloyd K.O. (2001) J. Biol. Chem. 276:27371-27375.
4. Kui Wong N., *et al.* (2003) J. Biol. Chem. 278:28619-28634.