

YBB818Ra01 100µg Vascular Endothelial Growth Factor Receptor 1 (VEGFR1) Organism Species: Rattus norvegicus (Rat)

> Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

> kDa 94 66.2 45 33 26 20 14.4 15% SDS-PAGE

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Ser303^{Val524} (Accession # P53767), with two N-terminal Tags, His-tag and S-tag. Host: E. coli Subcellular Location: Single-pass type I membrane protein. Cell membrane; Endosome. Purity: >95% Endotoxin Level: $\langle 1.0EU \text{ per } 1 \mu g \text{ (determined by the}$ LAL method). Formulation: Supplied as lyophilized form in PBS, pH7.4 containing 5% sucrose, 0.01% sarcosyl. Predicted isoelectric point:

The possible reasons that the actual band size differs from the predicted are as follows: 9.1 Predicted Molecular Mass: 30. 7kDa Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing

conditions. Applications: SDS-PAGE; WB; ELISA; IP.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

(May be suitable for use in other assays to be determined by the end user.)

Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

「 STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that incubate the protein at 37°C for 48h, and no obvious degradation and is. precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and S-tag, its sequence is listed below. MHHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF-SRDKGLYT CRVKSGSSFR TFNTSVHVYE KGFISVKHRK QQVQETIAGK RSHRLSMKVK



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com AFPSPEVVWL KDGVPATEKS ARYSVHGYSL IIKDVTAEDA GDYTILLGIK QSKLFRNLTA TLIVNVKPQI YEKSVSSLPS PPLYPLGSRQ VLTCTVYGIP QPTIKWLWHP CHYNHSKERN DFCFGSEESF ILDSSSNIGN RIEGITQRMM VIEGTNKTVS TLVV

[<u>REFERENCES</u>]

1. Brennan M., et al. (2012) Eur. J. Clin. Pharmacol. 68:645-655. 2. Huang H., et al. (2011) PLoS ONE 6:e21411-e21411. 3. Jesmin S., et al. (2012) Inflammation 35:484-500. 4. Yang K.Y., et al. (2011) Crit Care 15:R11-R11.