

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA618Hu01 100µg

Tachykinin Receptor 2 (TACR2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Val92~Leu290 (Accession # P21452), with N-

terminal His-Tag.

Host: E. coli

Subcellular Location: Cell membrane;

Multi-pass membrane protein.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS,

pH7.4, containing 5% sucrose, 0.01% sarcosyl.


Predicted isoelectric point: 9.1

Predicted Molecular Mass: 24.1kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MG H H H H H H S G SE F - VYA S H N I W Y FG R A F C Y F Q N LF P I TA M F V S IY S M TA I A A D

RYMAIVHPFQ PRLSAPSTKA VIAGIWLVAL ALASPQCFYS TVTMDQGATK CVVAWPEDSG

GKTLLLYHLV VIALIYFLPL AVMFVAYSVI GLTLWRRAVP GHQAHGANLR HLQAMKKFVK

TMVLVVLTFA ICWLPYHLYF ILGSFQEDIY CHKFIQQVYL

than 5% within the expiration date under appropriate storage condition.