

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB636Hu01 100µg

Recombinant Peptidylprolyl Isomerase C (PPIC)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Pro5~Val201

Tags: Two N-terminal Tags, His-tag and GST-tag

Accession: P45877

Host: E. coli

Subcellular Location: Cytoplasm.

Purity: >90%

Endotoxin Level: <1.0EU per 1 µ g (determined by the LAL

method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,

containing 5% trehalose, 0.01% sarcosyl.


Predicted isoelectric point: 8.9

Predicted Molecular Mass: 51.1kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80° C for 12 months. Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37° C for 48h, and no obvious degradation and

than 5% within the expiration date under appropriate storage condition.

is, incubate the protein at 37 C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less

[SEQUENCES]

The sequence of the target protein is listed below.

PRLLLP LVLCVGLGAL VFSSGAEGFR KRGPSVTAKV FFDVRIGDKD VGRIVIGLFG

KVVPKTVENF VALATGEKGY GYKGSKFHRV IKDFMIQGGD ITTGDGTGGV SIYGETFPDE

NFKLKHYGIG WVSMANAGPD TNGSQFFITL TKPTWLDGKH VVFGKVIDGM TVVHSIELQA

TDGHDRPLTN CSIINSGKID V