

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Visceral Adipose Tissue Derived Serine Protease Inhibitor (Vaspin) Organism: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[DESCRIPTION]

Protein Names: Visceral Adipose Tissue Derived

Serine Protease Inhibitor

Synonyms: Vaspin, Serpina12

Species: Mouse

Size: 100µg

Source: Escherichia coli-derived


Subcellular Location: Secreted.

[PROPERTIES]

Residues: Asn71~GIn148 (Accession # Q7TMF5), with N-

terminal His-Tag.

Grade & Purity: >95%, 12kDa as determined by SDS-PAGE reducing conditions.



15% SDS-PAGE

Formulation: Supplied as lyophilized form in PBS, pH 7.4,

containing 5% sucrose, 0.01% sarcosyl.

Endotoxin Level: <1.0 EU per 1µg (determined by

the LAL method).

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 10.5kDa



Predicted isoelectric point: 6.1



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSFF-NIFI SPI SIS TAFSMI SI GA QNSTI FFIRF GENEKEMSNW DVHAAFHYLL HKLNQETEDT KMNLGNALFM DQKLRPQQ