TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA009Mu01 50µg Recombinant Angiopoietin 2 (ANGPT2) Organism Species: Mus musculus (Mouse)

Instruction manual

kDa

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Tyr19~Ala147 (Accession # 015123),

with N-terminal His-Tag.

Host: E. coli

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in


PBS, pH7.4, containing 1mM DTT, 5% trehalose,

0.01% sarcosyl and preservative.

Predicted isoelectric point: 5.6

Predicted Molecular Mass: 16.4kDa

Applications: SDS-PAGE; WB; ELISA; IP.





227 Web:www.ybio.net Email:shybio@126.com

15% SDS-PAGE

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHHHSGSEF-YS NFRKSVDSTG RRQYQVQNGP CSYTFLLPET DSCRSSSSPY

MSNAVQRDAP LDYDDSVQRL QVLENILENN TQWLMKLENY IQDNMKKEMV EIQQNVVQNQ

TAVMIEIGTS LLNQTAAQTR KLTDVEA

[REFERENCES]

- 1. Maisonpierre P.C., et al. (1997) Science 277:55-60.
- 2. Carninci P., et al. (2005) Science 309:1559-1563.
- 3. The MGC Project Team. (2004) Genome Res. 14:2121-2127.
- 4. Yang W. J., et al. (2005) J. Biol. Chem. 280:9330-9335.