



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB572Hu01

50μg

Recombinant Lymphocyte Function Associated Antigen 1 Alpha (LFA1a)

Organism Species: Homo sapiens (Human)

***Instruction
manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

**9th Edition (Revised in Jul,
2013)**

[PROPERTIES]

**Residues: Asn809~His1066 (Accession # P20701),
with two N-terminal Tags, His-tag and T7-tag.**

Host: *E. coli*

**Subcellular Location: Membrane; Single-pass
type I membrane protein.**


Purity: >95%

**Endotoxin Level: <1.0EU per 1μg (determined by the
LAL method).**

**Formulation: Supplied as lyophilized form in PBS, pH7.4,
containing 5% sucrose, 0.01% sarcosyl.**

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 32.7kDa



The possible reasons that the actual band size differs from the predicted are as follows:

**Accurate Molecular Mass: 36kDa as determined by SDS-PAGE reducing
conditions.**

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.**
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.**



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGSSHHHHSSGLVPRGSHMASMTGGQQMGRGSEF-NLEEDAYWVQLD
LHFPPGLSFRKVEMLKPHSQIPVSCEELPEESRLLSRALS CNVSSPIFKA GHSVALQMMF
NTLVNSSWGD SVELHANVTC NNEDSDLLED NSATTIIPIL YPINILIQDQ EDSTLYVSFT
PKGPKIHQVK HMYQVRIQPS IHDHNIPTLE AVVGVPQPPS EGPITHQWSV QMEPPVPCHY
EDLERLPDAA EPCLPGALFR CPVVFRQEIL VQVIGTLELV GEIEASSMFS LCSSLISIFN
SSKHFFH

[REFERENCES]

1. Larson R.S., *et al.* (1989) J. Cell Biol. 108:703-712.
2. Nueda A., *et al.* (1993) J. Biol. Chem. 268:19305-19311.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

3. Cornwell R.D., *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:4221-4225.
4. Qu A., Leahy D.J. (1995) *Proc. Natl. Acad. Sci. U.S.A.* 92:10277-10281.