

YBB431MuO1 100µg Recombinant Mitogen Activated Protein Kinase 7 (MAPK7) Organism Species: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[PROPERTIES] kDa 70 44 Residues: Pro185^{Met419} (Accession # Q9WVS8), with two 33 N-terminal Tags, His-tag and T7-tag. 26 22 Host: E. coli 18 Subcellular Location: Cytoplasm. Nucleus. Nucleus, PML body. 14 Purity: >95% 10 Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). 15% SDS-PAGE Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl. Predicted isoelectric point: 6.2 Predicted Molecular Mass: 30.5kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

[<u>USAGE</u>]

9th Edition (Revised in Jul, 2013)

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Reconstitute in sterile PBS, pH7.2-pH7.4.

[<u>STORAGE AND STABILITY</u>]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below. MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- PSNLLV NENCELKIGD FGMARGLCTS PAEHQYFMTE YVATRWYRAP ELMLSLHEYT QAIDLWSVGC IFGEMLARRQ LFPGKNYVHQ LQLIMMVLGT PSPAVIQAVG AERVRAYIQS LPPRQPVPWE TVYPGADRQA LSLLGRMLRF EPSARISAAA ALRHPFLAKY HDPDDEPDCA PPFDFAFDRE ALTRERIKEA IVAEIEDFHA RREGIRQQIR FQPSLQPVAS EPVCPDVEM